

Orbactiv[®] & Kimyrsa[™] Referral Form

Please complete the following and send with demographics sheet, H&P, progress notes, medication list, and lab results to: p: 844.575.1515 | f: 844.797.5050 | e: specialtyreferrals@soleohealth.com

REFERRAL PROCESS	
1. PATIENT INFORMATION	2. PHYSICIAN INFORMATION
Name:	Physician's Name:
Address:	ARNP/PA:
City: State: Zip:	License #:
Home Phone: Mobile:	NPI#:
Caregiver Name:	DEA #: PA #:
Caregiver Relationship:	Hospital:
Caregiver Phone:	Office Contact:
Discharge Date: Start of Care Date:	Address:
Patient signed a DNR? Yes No	City: State: Zip:
Email:	Phone:
DOB: Social Security #:	Fax:
Gender: M F Nonbinary Height: Weight:	Nursing Agency:
Allergies:	Nursing Agency Phone:
3. DIAGNOSIS Please complete with ICD code and description.	
4. INSURANCE INFORMATION Please submit copies of the front and back of primary and secondary insurance cards with this referral.	
5. ORBACTIV or KIMYRSA ADMINISTRATION ORDERS* Carefully follow dose preparation and administration guidelines per manufacturer	
Orbactiv 1200 mg/1000 ML D5WIV over 3 hours x 1 dose, administered by a registered nurse	
☐ Orbactiv	
☐ Kimyrsa 1200 mg/250 ML NS or D5W IV over 1-hour x 1 dose, administered by a registered nurse	
Kimyrsa * Note: A clinical pharmacist from Soleo Health is available for consultation at the phone number at the top of this form.	
FLUSH ORDERS PIV/midline/PICC: Flush before, after each infusion, and as needed with 3-20 mL D5W (Orbactiv) or NS (Kimrysa), followed by Heparin 10	
unit/mL,2-5 mL if indicated Port: Flush before, after each infusion, and as needed with 5-20 mL D5W (Orbactiv) or NS (Kimrysa), followed by Heparin 100 unit/mL 5 mL.	
ANAPHYLAXIS ORDERS	
Adults: For mild reaction-administer diphenhydramine 50mg IVP/PO x 1. For severe reactions-administer epinephrine 0.3mg SubQ/IM x 1. May repeat x1 if needed. Contact 911 and monitor vital signs.	
Other: Labs:	
6. NURSING ORDERS: RN to assess, administer, and/or instruct patient/caregiver on medication, administration, vascular access device, disease process, signs or symptoms of adverse effects or complications of therapy. RN to establish, maintain and discontinue vascular access device. RN to draw labs as ordered via vascular access device or venipuncture.	
7. Dispense as written	
PHYSICIAN'S SIGNATURE (required):	Date: