

Our Mission

Fostering a culture of innovation.
Leading healthcare solutions.
Simplifying complex care.



www.soleohealth.com

Administration of IV Push Medications



ADMINISTRATION OF IV PUSH MEDICATIONS

Your medication has been pre-mixed by the Soleo Health pharmacist and placed in a plastic bag. Store all medication as directed. Some medications need to be refrigerated for stability.

- Check the label of each bag for your name, name of the medication, volume and the expiration of the drug.
- Each dose should be free of cracks, leaks, floating objects and discoloration. If you have any doubts, call your Soleo Health pharmacist.
- Remove one dose from your refrigerator one (1) hour before infusing to allow the medication to warm to room temperature. **NEVER HEAT OR MICROWAVE** the medication.
- Use each syringe only once.

SCAN THE QR CODE BELOW TO VIEW THE VIDEO



STEP 1

Wash hands thoroughly using soap and water for at least 20 seconds. Dry with a paper towel. Use your towel to turn off the faucet.

STEP 2

Clean your workplace and gather your supplies.

1. MEDICATION SYRINGE



2. ALCOHOL WIPES



3. FLUSH SUPPLIES

Flush Syringes



Antimicrobial Caps



STEP 3

Inspect your IV site for:

REDNESS

SWELLING

TENDERNESS

WARMTH

**IF THERE IS A CHANGE,
DO NOT INFUSE AND
CONTACT YOUR NURSE**

STEP 4

Inspect your medication.

IS IT EXPIRED?

IS YOUR NAME ON
THE LABEL?

IS IT THE RIGHT
MEDICATION?

ARE THERE ANY VISIBLE SPECKS OR FLOATING PARTICLES
IN THE MEDICATION?



STEP 5

Swab the end of your needleless connector using an alcohol wipe. Allow to air dry. Do not blot or fan.



STEP 6

Remove the cap from the saline flush. Holding the syringe upright, gently pull the plunger back to loosen it, and then push the plunger up to remove the air from the syringe. Attach saline flush syringe to catheter adapter by firmly pushing and twisting. Gently but firmly flush your catheter with ____ mL of saline.



STEP 11

Flush the IV catheter with saline flush solution and swab the end of your needleless connector using an alcohol wipe as before. If instructed, follow with a Heparin flush using units/mL ____ mL. Call your nurse if you cannot flush your catheter or meet resistance.



STEP 7

Swab the end of your needleless connector using an alcohol wipe. Allow to air dry. Do not blot or fan.



STEP 8

Attach the medication syringe to your needleless connector.



STEP 12

Attach antimicrobial cap to IV needleless connector, if provided.



STEP 9

Slowly push your medication over ____ minutes as directed. When finished, remove the medication syringe from your needleless connector.



STEP 13

Discard the used medication syringe in the appropriate waste container.



NOTES
